

NUMERICAL PROBLEMS

18.1 The length of a space ship is measured to be exactly one-third of its proper length. What is the speed of the spaceship relative to the observer?

Solution

Proper length = L_0

Measured length $L = \frac{L_0}{3}$ $v = ?$

Formula for Lorentz length contraction is

$$L = L_0 \sqrt{1 - \left(\frac{v^2}{c^2}\right)}$$

According to given condition ($L = \frac{L_0}{3}$) above equation can be written as

$$\frac{L_0}{3} = L_0 \sqrt{1 - \left(\frac{v^2}{c^2}\right)}$$

$$\Rightarrow \frac{1}{3} = \sqrt{1 - \left(\frac{v^2}{c^2}\right)} \quad \text{Squaring both sides \& solving for}$$

v

\Rightarrow

$$\frac{1}{9} = 1 - \left(\frac{v^2}{c^2}\right) \Rightarrow \frac{v^2}{c^2} = \frac{8}{9}$$

$$v = 0.9428 c$$

18.2 The time period of a pendulum is measured to be 3s in inertial frame of the pendulum. What is the period when measured by an observer moving with a speed of $0.95c$ with respect to the pendulum?

Solution

$t_0 = 3 \text{ sec}$

$t = ?$ If speed $v = 0.95 c$

Equation for time dilation is

$$t = \frac{t_0}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}} = \frac{3}{\sqrt{1 - \left(\frac{(0.95c)^2}{c^2}\right)}} = \frac{3}{\sqrt{1 - (0.9025)}}$$

$$t = 9.6 \text{ sec}$$

18.3 An electron, which has a mass 9.11×10^{-31} kg, moves with a speed of $0.75c$. Find its relativistic momentum and compare this value with the momentum calculated from classical expression.

Solution

$m_e = 9.11 \times 10^{-31}$ kg $v = 0.75 c$

Relativistic momentum $P_r = ?$ and compare with $P_{classical}$

$$P_r = m v = \frac{m_o v}{\sqrt{1 - \left(\frac{v}{c}\right)^2}}$$

$\frac{P_r}{P_{classical}} = 1.51$ $P_r = 1.51 P_{classical}$

$$P_r = \frac{(9.11 \times 10^{-31})(0.75 \times 3 \times 10^8)}{\sqrt{1 - \left(\frac{0.75 c}{c}\right)^2}}$$

$$P_r = 3.1 \times 10^{-22} \text{ kg m s}^{-1}$$

$$P_{classical} = m_o v$$

$$P_{classical} = (9.11 \times 10^{-31})(0.75 \times 3 \times 10^8)$$

$$P_{classical} = 2.05 \times 10^{-22} \text{ kg m s}^{-1}$$

For comparison, we find % difference as $\left(\frac{P_r - P_{classical}}{P_{classical}}\right) \times 100\% = 51\%$

18.4 An electron moves with a speed of $v = 0.85c$. Find its total energy and K.E in electron volt.

Solution

$v = 0.85c$

T.E = ? (in eV)

E = ? (in eV)

At the given speed ($0.85c$) the relativistic mass m is

$$m = \frac{m_o}{\sqrt{1 - \left(\frac{v}{c}\right)^2}} = \frac{9.11 \times 10^{-31}}{\sqrt{1 - \left(\frac{0.85 c}{c}\right)^2}} = 1.73 \times 10^{-30} \text{ kg}$$

Now Total energy T.E is

$$T.E = mc^2 = (1.73 \times 10^{-30})(3 \times 10^8)^2$$

$$T.E = 1.56 \times 10^{-13} \text{ J}$$

divide by $1.6 \times 10^{-19} \text{ J / eV}$

$$T.E = 0.973 \text{ MeV}$$

$$E_o = m_o c^2 = (9.11 \times 10^{-31})(3 \times 10^8)^2$$

$$E_o = 8.199 \times 10^{-31} \text{ J}$$

divide by "e"

$$E_o = 0.5124 \text{ MeV}$$

Now total energy T.E and kinetic energy K.E are related to rest energy E_o

$$T.E = K.E + E_o$$

$$K.E = T.E - E_o$$

$$K.E = (0.973 - 0.5124) \text{ MeV}$$

$$K.E = 0.461 \text{ MeV}$$

18.5 Rest mass of a proton is 1.67×10^{-27} kg. At what speed would the mass of the proton be tripled?

Solution

$m_o = 1.67 \times 10^{-27}$ kg $v = ?$ at which $m = 3m_o$

Relativistic mass of proton is given by

$$m = \frac{m_0}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}}$$

According to the given condition ($m = 3m_0$)

$$3m_0 = \frac{m_0}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}}$$

taking reciprocal

$$\frac{1}{9} = 1 - \left(\frac{v^2}{c^2}\right) \Rightarrow \frac{v^2}{c^2} = \frac{8}{9}$$

Taking square root

$$v = 0.9428c$$

18.6 At what fraction of speed of light must a particle move so that its K.E is one and a half times its rest energy?

Solution

$V = ?$

Total energy

when $K.E = \frac{3}{2} E_0 = \frac{3}{2} (m_0 c^2)$

$E =$ kinetic energy K.E + rest energy E_0

$$mc^2 = \frac{3}{2} (m_0 c^2) + (m_0 c^2) \quad \text{divide by } c^2$$

$$m = \frac{3}{2} m_0 + m_0 = \frac{5}{2} m_0$$

$$\frac{m_0}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}} = \frac{5}{2} m_0$$

$$\text{As } m = \frac{m_0}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}}$$

$$\frac{1}{\sqrt{1 - \left(\frac{v^2}{c^2}\right)}} = \frac{5}{2}$$

Take reciprocal and square

$$1 - \left(\frac{v^2}{c^2}\right) = \frac{4}{25}$$

$$\frac{v^2}{c^2} = \frac{21}{25}$$

Taking square root & solving for "v"

$$v = 0.916c$$

18.7 A metal, whose work function is 3.0 eV, is illuminated by light of wavelength $3 \times 10^{-7} \text{m}$. Calculate (a) The threshold frequency, (b) The maximum energy of photoelectrons (c) The stopping potential.

Solution

$$\Phi = 3 \text{eV} = 3 \times 1.6 \times 10^{-19} = 4.8 \times 10^{-19} \text{J}$$

$$\lambda = 3 \times 10^{-7} \text{m} \quad \text{(a) } f_0 = ?$$

$$\text{(b) } K.E_{\text{max}} = ?$$

$$\text{(c) } V_0 = ?$$

(a) Work function is the minimum amount of energy required for ejecting electrons from a metal surface

$$\Phi = h f_0$$

$$f_0 = \frac{\phi}{h} = \frac{4.8 \times 10^{-19}}{(6.63 \times 10^{-34})} = 0.72 \times 10^{15} \text{ Hz}$$

(b) According to the Einstein's equation of photoelectric effect

$$E = \Phi + K.E_{\max}$$

$$K.E_{\max} = hf - \phi$$

Since

$$f = \frac{c}{\lambda} = \frac{3 \times 10^8}{(3 \times 10^{-7})} = 10^{15} \text{ Hz}$$

$$K.E_{\max} = (6.63 \times 10^{-34})(10^{15}) - (4.8 \times 10^{-19})$$

$$K.E_{\max} = 1.83 \times 10^{-19} \text{ J} \quad \text{divide by } 1.6 \times 10^{-19}$$

$$K.E_{\max} = 1.14 \text{ eV}$$

(c)

$$K.E_{\max} = eV_0 \quad \text{where } V_0 \text{ is stopping potential}$$

⇒

$$V_0 = \frac{K.E_{\max}}{e} = \frac{1.14 \times 1.6 \times 10^{-19}}{1.6 \times 10^{-19}}$$

⇒

$$V_0 = 1.14 \text{ V}$$

18.8 The thermal radiation from the sun peaks in the visible part of the spectrum.

Estimate the temperature of the sun.

Solution

T = ? For visible part of spectrum

For visible part of spectrum, wavelength is 500 nm

$$\lambda_{\max} = 500 \text{ nm}$$

Using Wien's displacement law

$$\lambda_{\max} T = 0.2898 \times 10^{-2} \text{ m K}$$

$$T = \frac{0.2898 \times 10^{-2}}{\lambda_{\max}} = \frac{0.2898 \times 10^{-2}}{500 \times 10^{-9}}$$

$$T = 5796 \text{ K} \approx 5800 \text{ K}$$

18.9 A 50 keV X-ray is scattered through an angle of 90°. What is the energy of X-ray after Compton scattering?

Solution

Initially E = 50 keV = 50,000 × 1.6 × 10⁻¹⁹ J = 8 × 10⁻¹⁵ J

θ = 90° Energy after scattering E' = ?

Energy of X-ray photon after scattering can be found by equation

$$E' = \frac{hc}{\lambda'}$$

λ' is wavelength of scattered photon and is given by the equation (1)

$$\lambda' = \lambda + \frac{h}{m_0 c} (1 - \cos\theta) \quad (2)$$

λ is the only unknown at the right hand side for which we use

$$E = \frac{hc}{\lambda}$$

$$\lambda = \frac{hc}{E} = \frac{(6.63 \times 10^{-34})(3 \times 10^8)}{8 \times 10^{-15}}$$

λ = 2.49 × 10⁻¹¹ m putting this value in eq (2), we get;

$$\lambda' = 2.49 \times 10^{-11} + \frac{6.63 \times 10^{-34}}{(9.11 \times 10^{-31})(3 \times 10^8)} (1 - \cos 90)$$

$$\lambda' = 2.73 \times 10^{-11} \text{ m} \quad \text{put this in eq (1) we get}$$

$$E' = \frac{(6.63 \times 10^{-34})(3 \times 10^8)}{2.73 \times 10^{-11}}$$

$$E' = 7.28 \times 10^{-15} \text{ J}$$

$$E' = 45.5 \text{ keV}$$

divide by "e"

18.10 Calculate the wavelength of de Broglie waves associated with electrons accelerated through a potential difference of 200 V.

$\lambda = ?$ (for electron)

$$V_0 = 200 \text{ V}$$

De-Broglie's wavelength associated with a moving electron is given by

$$\lambda = \frac{h}{m v} \quad (1)$$

$$\text{Kinetic energy } \frac{1}{2} m v^2 = e V_0$$

$$v = \sqrt{\frac{2 e V_0}{m}} = \sqrt{\frac{2 (1.6 \times 10^{-19}) (200)}{9.11 \times 10^{-31}}}$$

$$v = 8.38 \times 10^6 \text{ m/s}$$

$$\text{Put this value in eq (1) we get } \lambda = \frac{6.63 \times 10^{-34}}{(9.11 \times 10^{-31})(8.38 \times 10^6)} = 0.87 \text{ \AA}$$

18.11 An electron is accelerated through a potential difference of 50V. Calculate its de Broglie Wavelength.

$$V_0 = 50 \text{ V}$$

$\lambda = ?$ (for electron)

Solution

De-Broglie's wavelength associated with a moving electron is given by

$$\lambda = \frac{h}{m v} \quad (1)$$

Since

$$\frac{1}{2} m v^2 = e V_0$$

Re-arranging them we get

$$v = \sqrt{\frac{2 e V_0}{m}}$$

$$v = \sqrt{\frac{2 (1.6 \times 10^{-19}) (50)}{9.11 \times 10^{-31}}} = 4.19 \times 10^6 \text{ m/s}$$

$$\lambda = \frac{6.63 \times 10^{-34}}{(9.11 \times 10^{-31})(4.19 \times 10^6)} = 1.74 \text{ \AA}$$

Put this value in eq (1) we get

18.12 The speed of an electron is measured to be $5 \times 10^3 \text{ m/s}$ to an accuracy of 0.003%. Find the uncertainty in determining the position of this electron.

Solution

$$v = 5 \times 10^3 \text{ m/s}$$

$$\text{Accuracy } A_c = 0.003\% = \frac{0.003}{100} = 3 \times 10^{-5}$$

Uncertainty in position $\Delta x = ?$ (for electron)

Uncertainty in velocity

$$\Delta v = 5 \times 10^3 \times A_c$$

$$\Delta v = 5 \times 10^3 \times (3 \times 10^{-5}) = 0.15 \text{ m/s}$$

Now according to Heisenberg's uncertainty principle,

$$\Delta x \Delta P = h$$

(As $\Delta P = m \Delta v$)

$$\Delta x = \frac{h}{m \Delta v} = \frac{6.63 \times 10^{-34}}{(9.11 \times 10^{-31})(0.15)} = 4.85 \times 10^{-3} \text{ m}$$

18.13 The life time of an electron in an excited state is about 10^{-8} sec. What is its uncertainty in energy during this time?

Solution

$$\Delta t = 10^{-8} \text{ sec}$$

$$\Delta E = ?$$

By Heisenberg's uncertainty principle,

$$\Delta E \Delta t = h$$

$$\Delta E = \frac{h}{\Delta t} = \frac{6.63 \times 10^{-34}}{10^{-8}} = 6.63 \times 10^{-26} \text{ J}$$