Mcqs no 02.. Pin point pupil are seen in A.alcohol poisoning B.morphine poisoning C.amphetamine poisoning D.cocaine poisoning E.LSD poisoning

5. The skull fracture that doesn't cross sutural lines is diastatic fracture

9. Synesthesia and flash backs occur in?(poison names)

13. Duration of detention for the purpose of assessment of mental health illness in kpk mental health Act 2017 is

A.28 days

B.22 days

C. 4 months

D.2 months

37. Gingival hyperplasia occurs with the use of which drug Answer : phenytoin

47. What are the advantages of double blinded study?

53.What is the most common mode of transmission of rabies to humans:

A ingestion of contaminated food

B mosquito bites

C inhalation of infected air

D contact with infected saliva through bites and scrapes

E sharing utensils with infected individuals

61. Q:Student t test measures

1:Difference bw mean

2:Difference bw 2 to 3 means

3:Difference bw frequency distribution

62. When there is no correlation between two variables it is true for

A\_ Negative correlation

B\_Null hypothesis (true)

70. What is complete transection of peripheral nerve called Neurotmesis

71. Most common cause of stroke worldwide is

A) Atherosclerosis

B) Hypertension

Don't remember other options

78. An infant born have bacterial meningitis

What is the most common cause of that

A E coli

B Streptococcus pneumonia

C Haemophilus influenza

D group b streptococci

80. At neuromuscular junction, motor neuron passes action potential to the muscle spindle fibre. The ion responsible for the release of ach from the synaptic vesicle is

- A. Calcium
- B. Nadium
- C. Potassium
- D. Chloride

86. The most common site of primary thrombosis causing cerebral infraction is a)middle cerebral artery

b) basillar artery

c)cavernous sinus

- d) carotid artery bifercation
- e) anterior cerebral artery

90: 45 years old lady presented with 3 days history of weakness of both lower limbs after 2 days . She has no previous history of hypertension and diabetes. 5 days back she had episode of diahrea resolve with treatment from local dr. Clinical examination shows flaccid paralysis of lower limbs with 1/5 power of lower limb and 2/5 power in upper limb. Planter are bilaterally going down. What is the diagnosis?

- a. poliomyelitis
- b. familial periodic paralysis
- c. acute hypokalimea due to Gi loss
- d. GB syndrome
- e. MS

92. 25 yrs old had bike accident 2 days ago presented with one day history of weakness of both lower limbs. Clinical examination shows bilateral up going plantar and sensory level up to umblicusWhich is most likely area of spinal cord injuryUpper dorsal spine

Lower dorsal spine

Upper lumbar spine

lower lumbar spine

Cauda equina

93) A 20 year old man presents with inability to hold his right hand straight due to an intramuscular injection in this side arm. Clinically he has right wrist drop. Which of the following explains his condition?

- A) Ulnar nerve injury
- B) Median nerve injury
- C) Radial nerve injury
- D) Extensor digitorum brevis injury
- E) Extensor digitorum longus injury

94. A 53 year old female with known rheumatoid arthritis has numbness and tingling sensation in her lateral three fingers of right hand there is sharp burning pain extending from right hand to right arm.

diagnose the case.

a. Ulnar nerve entrapment

- b.median nerve entrapment
- c. radial nerve entrapment
- d. Tenosynovitis
- e wrist joint subluxation

97. A 83 year old man with irregular talking and incontinence of urine for past 3 years. What is ur diagnosis.

A. Dementia

- B. Delirium
- C. Parkonson
- D.
- Ε.

101. Which type of meningitis is it ? Don't remember the scenario

- 1. Bacterial
- 2. Viral
- 3. Tuberculous

103.A 2 year old child presented with cerebral palsy.He had ataxic gait.How would you classify this ataxia.

a)acute ataxia

b)chronic non progressive ataxia

c)chronic progressive ataxia

d) intermittent ataxia

e)paroxysmal ataxia

Ans.b

118. The first you should do after bitten by a dog:

- 1. Go to hakeem
- 2. pain killer
- 3. apply tumeric
- 4. get anti rabies vaccine

5. wash wound with mild soap and running water for 10 minutes and then go to hospital