

# Introduction and classification of glaucoma and Congenital Glaucoma

Associate Professor Dr Afzal  
Qadir

Pediatric ophthalmologist  
Eye unit HMC


# Acknowledgement

- Becker- Schaffer's Diagnosis and therapy of The Glaucomas (8<sup>th</sup> Edition).
- Kanski's Clinical Ophthalmology (8<sup>th</sup> Edition).
- Comprehensive Ophthalmology (A.K.Khurana) (7<sup>th</sup> Edition).

# Learning Objectives

- **At the end of this class the students shall be able to :**
- Define and classify glaucoma.
- Define congenital glaucoma.
- Understand the pathogenesis and clinical features of congenital glaucoma.
- Understand the fundamentals of managing congenital glaucoma.

**Excessive  
blinking +/-  
watering**



**Hazy cornea  
Large corneas**

# Question

- A child presents with watering , photophobia and an enlarged cornea with a diameter of 13mm. Examination of the eye reveals double contoured opacities concentric to the limbus. Which of the following is the most likely diagnosis:
  - Superficial keratitis
  - Deep keratitis
  - Thyroid eye disease
  - Congenital glaucoma

# What is glaucoma ?


- The term glaucoma is derived from the Greek word “glaukos” meaning “gray blue”
- Second leading cause of blindness worldwide
- Third most common cause of blindness in India
- Not reversible



# Definition of glaucoma

- Group of disorders characterized by progressive optic neuropathy resulting in characteristic morphological changes at the optic disc leading to a specific pattern of irreversible visual field defects (with or without a raised IOP).

# Classification of glaucoma





# Primary glaucoma

- Open angle glaucoma



- Primary Open angle glaucoma
- Normal Tension glaucoma
- Juvenile Open angle glaucoma
- Secondary Open angle glaucoma



- Steroid induced glaucoma
- Pigmentary glaucoma

# Primary glaucoma

- Angle closure glaucoma



- Primary angle closure glaucoma
- Secondary angle closure glaucoma



- Swollen lens
  - Posterior segment tumours
  - Neovascular glaucoma
- 
- Plateau iris syndrome

# Childhood glaucoma

- Primary congenital glaucoma
- Glaucoma associated with ocular abnormalities
- Glaucoma associated with systemic abnormalities

# Secondary glaucoma

Glaucomas after ocular surgery

Steroid induced glaucoma

Traumatic glaucoma

# Childhood glaucoma-Introduction

- Diverse group of disorders
- Primary congenital glaucoma-

Developmental abnormality of angle of anterior chamber leading to high intraocular pressure(IOP).

- Secondary congenital glaucoma

With associated ocular and systemic anomalies

# **ANGLE OF ANTERIOR CHAMBER**

- The **peripheral recess** of anterior chamber is known as the angle of anterior chamber.
- It is clinically visualized by **gonioscopy**.
- Starting at the root of iris & progressing anteriorly towards the cornea, the following structures can be identified in a normal angle in an adult :
  - 1) Ciliary body band (CBB) & root of iris
  - 2) Scleral spur (SS)
  - 3) Trabecular meshwork (TM)
  - 4) Schwalbe's line (SL)

# Angle of anterior chamber as seen on gonioscopy



*Structures visible on gonioscopy*


From Schwalbe's line to ciliary body

From Schwalbe's line to scleral spur



From Schwalbe's line to trabecular meshwork

Schwalbe's line only

None of the angle structures visible







# Childhood glaucomas

- **True congenital** glaucomas- At birth or during intrauterine period.
- **Infantile** glaucoma- Upto three years of age.
- **Juvenile** glaucoma- After three years of age and upto 35 years of age.

# Prevalence and genetic pattern

- Sporadic occurrence in most cases (90%)
- Autosomal recessive in 10% of cases
- Loci linked with congenital glaucoma are  
2p21(GLC3A), 1p36(GLC3B) and 14q24(GLC3C)
- 60% diagnosed by the age of 6 months and 80%  
diagnosed within the first year of life

# Prevalence and genetic pattern

- Bilateral (about 70%) but asymmetrical
- Boys are affected slightly more frequently than girls (65%)
- Prevalence is 1 in 10,000 births
- Chance of second sibling having disease is 3%
- Chance of third sibling (of two affected siblings) having disease is 25%

# Pathogenesis

- Faulty development of angle of anterior chamber from neural crest derived cells (**trabeculodysgenesis**)
- Absence of angle recess with flat/concave iris insertion.



- Impaired aqueous outflow



- Elevated IOP



The normal chamber angle: on the left is a histological cross-section; on the right is a drawing of the same



An underdeveloped chamber angle

# Clinical presentation

## Classic triad of



- Epiphora
- Blepharospasm
- Photophobia



- Babies rub their eyes
- Enlarged eyes
- Vision impaired


# Corneal signs

- Corneal oedema
- Corneal enlargement (Corneal diameter > 13mm)
- Haab's striae : Descemet's membrane is not very elastic and stretching may result in small linear/circumferential tears that cause a certain degree of corneal opacification.



# Clinical presentation

- **Buphthalmos**: Enlargement of the globe as a result of elevated IOP. All segments of the outer eye especially the cornea and **sclera** expand principally at the corneoscleral junction
- The anatomic landmarks are displaced.
- The anterior chamber is deep



Advanced developmental glaucoma with extensive enlargement and scarring of the cornea.




# Clinical presentation

- Sclera becomes thin and appears blue (due to underlying uveal tissue)
- Iris- atrophic in later stages
- Optic disc- variable cupping
- Intraocular pressure(IOP)- raised
- Axial myopia- due to increased axial length of eyeball


# Examination under anaesthesia

- **Mandatory in all cases**
- Includes :
- **Measurement of IOP –**  
Perkins tonometer/Tonopen  
(Normal 10-21 mm Hg)
- **Measurement of corneal diameter – by**  
callipers  
(Normal 9.5mm-10.5mm)



# Examination under anaesthesia

- Slit lamp examination- with portable slit lamp
- Ophthalmoscopy- to evaluate optic disc



Asymmetric disc cupping in a child with developmental glaucoma. (A) Note steep-walled cup. This is typical of glaucomatous cupping in the elastic infant eye. (B) The left eye has no cupping.

# Examination under anaesthesia

- **Direct Gonioscopy** – to examine angle of anterior chamber
- Koeppe's gonioscopy lens is preferable
- Angle is open but immature in congenital glaucoma



# Differential Diagnosis

- **Hazy/Cloudy cornea----**
- STUMPED (Sclerocornea, Trauma, Ulcer, Metabolic disorders, Peter's anomaly, Endothelial dystrophy)
- **Watering and intolerance to light-----**  
Congenital Naso Lacrimal Duct obstruction  
keratitis, conjunctivitis
- **Optic cupping ----** disc coloboma, hypoplasia, physiological cupping
- **Corneal enlargement --** megalocornea, high myopia
- **Descemet's breaks ---** Forceps delivery ,birth trauma

# Management

- **Glaucoma surgery** is the **primary option**
- Medications are not very effective
- Role of medical management is temporary, till surgery is taken up.
- Beta blockers (Timolol), hyperosmotic agents(Mannitol), carbonic anhydrase inhibitors (acetazolamide/dorzolamide)
- **Miotics and Alpha-2 agonists are not used in children.**

Goniotomy/Trabeculotomy

```
graph TD; A[Goniotomy/Trabeculotomy] --> B[Trabeculectomy with trabeculotomy]; B --> C[Modified Trabeculectomy]; C --> D[Glaucoma drainage implant]; D --> E[Cyclodestructive procedures];
```

A vertical flowchart with five colored rectangular boxes connected by downward-pointing arrows. The boxes are: 1. Red: Goniotomy/Trabeculotomy; 2. Green: Trabeculectomy with trabeculotomy; 3. Purple: Modified Trabeculectomy; 4. Blue: Glaucoma drainage implant; 5. Orange: Cyclodestructive procedures.

Trabeculectomy with  
trabeculotomy

Modified Trabeculectomy

Glaucoma drainage implant

Cyclodestructive procedures

# Approach to management

Goniotomy/Trabeculectomy/Combined Trabe-Trab

Surgical outcome?

EUA after 3-4 weeks

**IOP controlled**

Evaluation after 3 months

Normal IOP

Evaluation after 3 months

FU every 3 months

Record IOP, CDR, VA  
Axial length, VF (if possible)

VISUAL  
REHABILITATION

**IOP not controlled**

Add medical therapy

If IOP not controlled

**repeat Trab ± MMC**

Controlled

Uncontrolled


**Poor prognosis**

Consider  
**Drainage implant**  
**Cyclodestruction**<sup>32</sup>




# Goniotomy

- Safe procedure when performed skilfully.
- Performed with direct visualization of trabecular meshwork
- Aims to transect Schlemm's canal by ab-interno approach
- Incises only superficial trabecular tissues, necessary to cure this disease



# Trabeculotomy

- Ab-externo trabeculotomy has good success rates.




Trabeculotome


# Trabeculotomy with trabeculectomy

- Most commonly performed surgery in India
- Easy adaptability
- Safe and successful
- Suitable in compromised corneas
- More predictable results


# Steps of Trabeculectomy with trabeculotomy




Scleral flap



Dissection upto grey limbus



Trabecular meshwork cut



Diffuse subconjunctival bleb

# Role of antimetabolites in paediatric glaucoma

- Significantly **more complications** associated with the use of Mitomycin(**MMC**) in paediatric glaucomas
- Thin, avascular filtering blebs
- Wound leakage
- Choroidal detachment
- Bleb related endophthalmitis



# Options for refractory glaucoma ?

- Glaucoma Drainage Devices
- Cyclo-destruction

# What is a Glaucoma Drainage Device?

Glaucoma drainage devices (GDDs) create an alternate aqueous pathway from the anterior chamber (AC) by channeling aqueous out of the eye through a tube to a subconjunctival bleb. This tube is usually connected to an equatorial plate under the conjunctiva.




# Cyclodestructive procedures

- Cyclocryotherapy
- Cyclophotocoagulation
  - Transscleral
  - Transpupillary
  - Endoscopic



# CYCLO CRYOTHERAPY


TREATMENT OF 1950  
BIETTI



# Lasers relatively safer energy

- Trans-scleral route
- Direct application to ciliary epithelium

Trans pupillary



# Trans-scleral route diode laser

810 nm wave length


Penetrates through sclera

Contact delivery through  
fibre optic cable

Diode laser is preferred

Melanin in the ciliary  
epithelium better absorbs  
this wavelength

Causes more **targeted  
destruction** with  
less inflammation



# VISUAL REHABILITATION

- Correction of refractive error
- Management of media opacities
- Amblyopia therapy to achieve binocular stereoscopic vision



# VISUAL REHABILITATION

- Low vision aids
  - ❖ Telescopes (hand-held or spectacle-mounted)
  - ❖ Hand or pocket magnifiers (2× to 3×)




III. Hand held magnifiers

# CONCLUSIONS

- Glaucoma is a group of disorders characterized by progressive optic neuropathy.
- Early diagnosis and prompt treatment can preserve vision.
- All children with suspected childhood glaucoma should be examined under anaesthesia.
- Mainstay of management of childhood glaucoma is surgery
- Visual rehabilitation and counseling of the parents of the child is as important as IOP control.

# Question



- Identify the abnormality marked by arrow.
- Which structure is involved?
- What type of slit lamp illumination is used in the photograph?
- Mention one differential diagnosis of the condition





Thank you