



## **OSCE**

For Final Year MBBS NLE (Clinical Skills Examinations)

### 2nd Edition

**Covers OSCE Stations of Final Year Subjects** 


History Taking and Differential Diagnosis

Examinations

Instruments, X-Rays & ECGs

Procedures, Counselling, EPI

DR FAISAL HASSAN Dr Abdul Basit Dr Bushra Yousaf



#### Salient Features of Hassan's OSCE:

- · Covers Stations of Medicine, Surgery, Gynae / Obs & Paeds
- · History Taking and Differential Diagnosis
- Examinations
- Fluids Analysis
- · ABGs, Drugs and Poisoning
- Instruments
- X-Rays & ECGs
- Procedures
- Counselling
- · EPI





PKR Rs: 795/-

For Delivery All Over Pakistan © 0346-9049087 曾 0946-726817

## Hassan's

## **OSCE**

## For Final Year MBBS, NLE-2 2nd Edition

### **Note:**

- ✓ 2<sup>nd</sup> Edition is the Updated Edition in Best Quality Page and Completely Coloured Book
- Covers OSCE Stations of All Final Year Subjects
- ✓ The Most Authentic, Best Selling Book According to Pattern of KMU Exam
- ✓ History Taking and D/Ds
- ✓ All Systems Examinations
- **✓ Fluids Analysis**
- ABGs, Drugs and Poisoning
- ✓ Instruments
- ✓ X-Rays and ECGS
- ✓ Procedures
- ✓ Counselling
- ✓ EPI Schedule

# Published By: Swat Book House & Publications

For All Kind of Books, Medical Instruments, Overalls etc Whatsapp# 0346-9049087 Phone# 0946-726817

AVAILABLE IN ALL MEDICAL BOOKSHOPS OF KP, ISLAMABAD, PUNJAB & KARACHI

#### Note:

- ✓ In this pdf file Only Highlights of Book given.
- **✓** Total Number of Pages=353

#### **PREFACE**

When I was a student of final year MBBS at KMC, I found it very difficult to study for TOACs examination. Because one has to study different books for different OSCE stations. So I felt the need to combine all the relevant OSCE stuff in one book form, i.e. examination, instruments, and other stations.

Great effort was made in writing and compiling of this book. Most of the stuff is taken from standard books along with ward lectures and examination books.

This book almost covers all stations in medicine and Paediatrics TOACs. However, it should be kept in mind that without proper practice of these stations and without performing these examinations on patients, this book will give you theoretical clue.

Please let us know your feedback and suggestions for improvement of stuff in the next edition at

E-mail: kmcite152@gmail.com

**FACEBOOK PAGE: HASSAN OSCE GUIDE** 

Whatsapp#: 0345-3435070

DR. FAISAL HASSAN KMC PESHAWAR

#### THIS BOOK INCLUDES:

#### Ward lectures of

- DR. HUMAYUN (HOD MEDICAL WARDS)
- DR. HASHIM (PROF.MDW KTH)
- DR. AWAIS (AP. MDW KTH)
- DR. GHULAM SHABBIR (PROF. MCW KTH)
- DR. SALEEM IQBAL (PROF MBW KTH)
- DR. ZAHID FIDA (PROF. MEW KTH)
- DR. IRSHAD (HOD PEADS KTH)
- DR. QAMAR (PROF.PEADS KTH)
- DR. ZIA (PROF.PEADS KTH)
- DR. SABIR (PROF. PEADS KTH)
- DR.SABAHAT (AP PEADS KTH)

#### **POINTS FROM:**

- NICHOLAS TALLEY
- MACLOED'S
- BEDSIDE TECHNIQUES Etc.

#### **ACKNOWLEDMENT**

First of all I am extremely grateful to **My Family** for trusting and believing in me. Without whom I can't even think of myself.

Special thanks to **Doctor SANA** (TMO MAW KTH) who thoroughly checked every word one by one and pointed out the short comings. Once again thank you so much ma'am for your precious time, you really helped me a lot. (1st edition)

SPECIAL THANKS to **DR. ABDUL BASIT**, my most exacting critic, my closest friend without whom this book would not have been this book.

Special thanks to **Dr. BUSHRA YOUSAF** who helped me a lot in writing the Procedures and reviewing the Book.

Thanks to **NIDA ALAM** pharmacist for helping me a lot in compiling the book.

Gratitude to **Dr. HALEEMA WAZIR**, my junior who compiled the newly added stations in the TOACs.

Thanks to the **Dr. AZHAR HAYAT, Dr. ZOHAIB, Dr. AMMAR ZAHID, Dr. NASEEB, and Dr. SABIR** who always treated me as younger brother and encouraged me at every step.

Thanks to all the Contributors.

#### **CONTRIBUTORS**

I would like to give credit to these people for helping me in writing and collecting stuff.

- 1. Dr Abdul Basit
- **2.** Dr. Bushra Yousaf
- 3. Dr Haleema Wazir
- 4. Dr Ambareen Khan
- **5.** Dr Marhaba iqbal
- 6. Dr Syeda Kainat
- 7. Dr Saima Rahman
- 8. Dr Azhar Hayat
- 9. Dr Sabir
- **10.** Dr Rehan Haseen
- 11. Dr Naqeeb Ansari
- 12. Dr Fawad Nasir
- 13. Dr. Raza ul Haq
- 14. Dr Naseeb

## Page-9

### **INDEX**

SECTION: 1	XAMINATIONS
PAGE	
History taking & Differentials	5 11
Diagnosis	· · · · · · · · · · · · · · · · · · ·
General Physical Examinatio	n 34
Respiratory System Examina (Chest)	tion 58
Abdominal Examination (GIT	71
Diabetic Foot Examination & Thyroid	84
CVS Examination (precordium	n) 101
CNS Examination	122

SECTION: 2	FLUID ANALYSIS
CSF Analysis	160
Ascitic Fluid	167
Pleuritic Fluid	173
Urinalysis	175

SECTION: 3	ABGs, DRUGS &
POISONING	
	(0.5)

ABGs	180
Drugs	185
Poisoning	203

SECTION: 4	INSTRUMENTS, X-RAYS & ECG

1 Instruments	209
7 X-rays	247

Hassan	'c	ns	CE
паээан	15	US	CE.

: ECG

and	$\mathbf{r}_{\mathbf{J}}$		
2nd	Ła	III	on

272

Page-11

SECTION: 5	PROCEDURES, COUNSELLING, EPI,
INSTRUMENTS	

	3
Nebulization , Resuscitation, I/V	302
cannula, NG tube, catheter, CPR	
Counselling Steps	330
EPI schedule + Indications for Nursery admission	335
Gynae & Surgery Instruments	338-
	353
11253115	

### **HISTORY Taking**

"Always listen to the patient that might be telling you the diagnosis" (Sir William Osler)

First of all follow the basic steps

- Approach to the patient in friendly, sympathetic and gentle way
- Greeting + introduce yourself to the patient.
- Take consent + pay full attention to the patient.
- If the patient is eating or doing something (leaving for washroom etc.) wait for the patient or visit later.
- Use the patient own words and don't use difficult medical terminologies.
- First listen to the patient and don't interrupt the patient by asking leading questions.
- At the end you can ask the necessary questions if any to complete the History.

Record the history in the following pattern.

- 1. Personal profile
- **2.** Chief/presenting complaints (C/C)
- **3.** History of presenting illness (HOPI)
- 4. Review of systems (ROS)
- 5. Current medications and Drug history
- 6. Past medical/surgical History

- 7. Personal History
- 8. Family History
- 9. Allergy history
- 10. Travel history
- 11. Occupational History
- 12. Menstrual History (Female)
- 13. Socioeconomic history

#### 1. **PERSONAL PROFILE:**

- a) Name: (for identification of the patient say for example XYZ Son/daughter/wife of ABC.it is also possible that in the same ward/chamber two different patients with same name are there. So also ask for the father/husband name for differentiation)
- b) Age: (age is much more important because some diseases are more common in certain age group. For example stroke, malignancy, MI, etc. are more common in older group and measles, chicken pox etc. are more common in children)
- Sex: (some diseases are common in males (myocardial infarction) and some are present in females mostly i.e. SLE, primary biliary cirrhosis)
- d) **Address**: (for further communication and some diseases are more common in one place than the other)
- e) Religion:

- f) Marital status/Habitus: (unmarried people are mostly in depression, married/habitual may have some sexually transmitted disease.)
- g) **Occupation:** (many diseases are commonly related to occupation, and to know about the socioeconomic history of the patient)
- h) **Mode of Admission:** admitted through emergency or Pod. Also indicate about the nature of the disease.
- i) Date and Time

#### 2. CHIEF/PRESENTING COMPLAINTS:

The complaints which bring the patient to the doctor.

Recorded in chronological order i.e. the symptom started first will be written first and so on.

And along the symptom duration is also mentioned.

For Example

- 1. Abdominal pain...... 12 days
- **2.** Fever...... 10 days
- 3. Nausea / vomiting ...... 6 days

Note: use patient own words, no leading questions, recurrent symptoms can be noted

#### **HISTORY OF PRESENTING ILLNESS (HOPI)**

That part of the history in which the chief complaints are elaborated in detail.

It is actually the chronological description of present illness from initial sign to the present.

#### Elements of HOPI are:

- 1. Location/site
- **2.** Quality
- **3.** Severity
- 4. Duration
- **5.** Timing
- **6.** Modifying factors
- **7.** Associated sign and symptoms

#### For Example:

PAIN (SOCRATES)	FEVER (ODIPARA)
<b>S</b> = SITE	<b>0</b> = onset
<b>O</b> =ONSET	<b>D</b> = Duration
<b>C</b> = CHARACTER	I = intensity
(Sharp/dull,	(Low Grade or High Grade/
Heaviness, Aching,	Rigors and chills)
throbbing, colicky,	
Gripping,	

Burning/tingling,	
boring/ stabbing,	
Crushing,	
or doming,	
$\mathbf{R} = \text{RADIATION}$	P = pattern
	Continuous/intermittent/
	remittent/ hectic/ relapsing
	remittent/ nectic/ relapsing
$\mathbf{A} = ASSOCIATED$	<b>A</b> = Aggravating factors
SYMPTOMS	
T = TIMING	<b>R</b> = Relieving factors
(DURATION,	
COURSE, PATTERN)	(, P <sub>k</sub>
, , , , , , , , , , , , , , , , , , , ,	
<b>E</b> = EXACERBATING	<b>A</b> = Associated factors
& RELIEVING	
FACTORS	
S = SEVERITY	

### **DIFFERENTIAL DIAGNOSIS:**

#### **COUGH:**

ACUTE COUGH	CHRONIC COUGH
Inhaled foreign	Nonproductive:
Body	
Dody	1. Asthma
	2. Allergy
	3. Vocal Cord dysfunction
	4. COVID
	5. GERD
	6. Drug related like ACE
	inhibitors
	7. Post nasal drip
Respiratory tract	Productive:
infection	O'
	1. Pneumonia , Sinusitis ,
	Bronchitis
7,2	2. TB, lung cancer,
	pulmonary
200	embolism(all are blood
1000	stained)
17.0	3. COPD
	(purulent/mucoid)
	4. Bronchiectasis
	(purulent)
	<ol><li>Pulmonary edema</li></ol>
	(pink/frothy)

#### **ABDOMINAL PAIN**

GASTROINTESTINAL + PERITONEUM	GENITOURINARY + GYNAECOLOGICAL	Ascular Page 118
<ol> <li>PUD</li> <li>Gastritis</li> <li>Malignancy</li> <li>Volvulus</li> <li>Appendicitis</li> <li>Obstruction</li> <li>IBD</li> <li>Strangulated Hernia</li> <li>Intussusception</li> <li>Cholecystitis</li> <li>Pancreatitis</li> <li>Cholangitis/Hepatitis</li> <li>Diverticulitis</li> <li>Splenic Rupture</li> <li>Peritonitis</li> </ol>	1. UTI 2. Urinary Retention 3. Ureteric Colic 4. Polycystic Kidney Disease 5. Pyonephrosis 6. Ruptured Ectopic Pregnancy 7. Ovarian Cyst Torsion/Rupture 8. Severe Dysmenorrhea 9. Salinities 10. Endometritis 11. Fibroid Red Degeneration	5) Embolus
	etc	

#### **EXAMINATIONS**

- GENERAL PHYSICAL EXAMINATION
- RESPIRATORY SYSTEM EXAMINATION
- ABDOMINAL EXAMINATION
- DIABETIC FOOT EXAMINATION
- THYROID EXAMINATION
- CVS EXAMINATION
- CNS EXAMINATION

#### **GENERAL PHYSICAL EXAMINATION:**

First of all, follow the basic steps "RIPE"

R=Right side approach

I =Introduction and consent

P = Position (supine + arm abduction)

E=Exposure and Explain the procedure to the patient

#### **FIRST IMPRESSION OR GENERAL APPEARANCE:**

First impression of the patient's condition must be deliberately sought, whether patient feels well, mildly ill or severely ill.

#### **POSTURE AND ATTITUDE:**

It gives information about patient illness e.g.

 A pt. with severe heart failure prefers to sit propped up because his dyspnea worsens on lying flat

- A pt. with peritonitis lie down on the back, with two legs bending. (pain and cannot move)
- Squatting position in case of congenital heart disease.
- Severe colicky pain =restless
- In meningitis= neck retraction etc.

#### **CONSCIOUNESS:**

Note that pt. is well oriented in time, place and person or alert, confused and drowsy or unconsciousness by assessing level of unconscious by using Glasgow Coma Scale (GCS).

Behaviour	Response	5
	4. Spontaneously	300
	3. To speech	2
	2. To pain	
	1. No response	
Eye Opening Response		
	5. Oriented to time, person and place	1
	4. Confused	
	Inappropriate words	
	2. Incomprehensible sounds	
	1. No response	
Verbal Response		
- //	6. Obeys command	
	5. Moves to localised pain	
(P)	4. Flex to withdraw from pain	
	3. Abnormal flexion	
3	2. Abnormal extension	
Motor Response	No response	

- For Example: Brain injury is classified as on GCS
  - Severe= GCS < 8-9</li>
  - Moderate = GCS, 9-12
  - Mild = GCS > 13 or equal to 13

#### **PHYSIQUE:**

Visual impression is made about a pt. height and weight. Both should be measured and compared with ideal body weight and height. Compare the sitting height with arm span and total height.

Page-21

- **TALL STATURE:** Genetic, Marfan's syndrome, klinefelter syndrome etc.
- **SHORT STATURE:** constitutional, genetic, achondroplasia, cretinism, hypopituitarism
- In achondroplasia arms and legs are short while trunk is normal, so sitting height is more than legs.
- In Marfan's syndrome and hypogonadism arm span is more than double the sitting height.

#### **HANDS**:

Many pathologies can be detected from the hands of pt.

Shaking hands with pt. is important to know about the disease like dystrophic myotonia in which patient is unable to let go.

#### **SHAPE:**

- Hands adopt special shape in tetany due to carpel spasm
- Short 4<sup>th</sup> metacarpal in females is seen in Turner's syndrome.

#### SIZE:

Hands are large and broad in acromegaly

#### **NAILS:**

Blue nails: cyanosis, Wilson's disease, onchronosis

- **Red nails:** polycythemia, CO poisoning, sometimes normal finding
- **Yellow nails:** yellow nail syndrome, fungal infection, diabetes etc.
- **Brown nails:** Chronic kidney disease
- White nails with darker rim: Hepatitis
- Pale nails: Anemia, CHF, liver disease
- **Cracked or split nails**: Thyroid disease
- Brittle nails: (which can break easily) iron deficiency anemia, PVD, hypocalcaemia etc.

#### **CLUBBING:** (Hippocratic nails/fingers)

It is a physical sign characterized by bulbous enlargement of soft parts of terminal phalanges with both transverse and longitudinal curving of the nail.

Also called **watch glass nails**, drumstick fingers, Hippocratic nails

#### **TYPES:**

Acute: erythematous

**Chronic**: non erythematous




Figure: Normal Schamroth sign



Figure: Clubbing

## RESPIRATORY SYSTEM EXAMINATION OR CHEST EXAMINATION

## <u>COMMAND: EXAMINE FROM THE FRONT FOR CHEST OR RESPIRATORY SYSTEM</u>

First of all, follow the basic steps "RIPE"

R=Right side approach

I =Introduction and consent

P = Position (supine + arm abducted)

E=Exposure & explain the procedure.

#### **Exposure:**

**Take off the shirt in male** and maintain privacy in females or do it in presence of female colleague or nurse **Position:** 

While examining from the front patient should lie supine with arms abducted or patient lying at an angle of 45.

#### **INSPECTION**

Inspect the front and sides from the foot end. Note for the following

- Shape and symmetry of chest
- Look at the face for SOB, cough and cyanosis
- Respiratory rate counting (missed by most of the students)
- Type of respiration
- Prominent veins
- Any scars
- Deformity
  - Pectus carinatum (pigeon chest)

- Pectus excavatum (funnel chest)
- Harrison's sulcus
- Any striae
- Note any wheezing
- Flail rib
- Any mass
- Chest recession or retractions
- Observe the surrounding i.e. any nebulizer, medicine etc. (accessory equipment)

#### **RESPIRATORY RATE:**

Count the respiratory rate for one complete minute (IMP)

Age	Normal RR (breaths/min)	Tachypneic (breaths/min)
0-2 months	Up to 59	60 or >60
2 months - 1 year	Up to 49	50 or > 50
1-5 years	Up to 39	40 or > 40
> 5 years	15-20	>20

NOTE: count the RR in inspection as it is noted by examiner especially in peads.

#### **MOVEMENT OF CHEST:**

- Reduce movements in:
  - 1) Pleural effusion
  - 2) Collapse, fibrosis
  - 3) Consolidation
- Abnormal Movements: airway obstruction i.e. Emphysema, asthma etc.

#### **TYPES OF BREATHING:**

- Thoracoabdominal: normal in females
- Abdominothoracic: normal in males and children
- Periodic breathing: in babies (breath fast several times and then have a brief rest for less than 10 sec.)
- Acidotic breathing: renal failure and ketoacidosis
- Cheyne-stokes breathing: over ventilation alternates with complete apnea i.e. LVF, Inc ICP, narcotic overdose

#### **PALPATION**

While starting palpation

- 1) Warm your hands.
- 2) Ask about pain and tenderness (missed by most of the students).

#### 1. PALPATION OF TRACHEA:

The patient should lie straight. Trachea can be palpated in the following two position.

- 1) Either remove the pillow from the pt. head
- 2) Ask the pt. to extend the neck and look backwards or toward the roof.

Trachea can be palpated by one, two or three finger method.

Ideally, we use three finger method to note any deviation of trachea.

Stabilize middle finger in midline and put other fingers around the midline in space and note any deviation as shown in the fig.



Normally trachea is somewhat towards the right side but not so much palpable....we consider it in the center as normal.

Conditions which:

- Pull the trachea: collapse, fibrosis, agenesis of lungs etc.
- **Push the trachea**: pneumothorax, pleural effusion (Air+ water), Kyphoscoliosis.

#### 2. CHEST EXPANSION:

Grasp the sides of the chest with the fingers in such a way that the thumbs of both hands approximate in the midline as shown below. Now ask for deep breath and note any abnormality or symmetry.



Causes of Reduced chest movements:
UNILATERAL BILATERAL

Pleural effusion and pneumothorax	Emphysema
Chest trauma	Hydrothorax
Fibrosis of lung	Bronchial asthma
Consolidation	Myasthenia gravis
Chest trauma	Obesity

After palpation with hands you can also check and measure the chest expansion with measuring tape.

#### 3. VOCAL FREMITUS:

Place the edge of hand on the chest horizontally and ask the patient to say 1, 2, and 3 and compare both sides.

NOTE: examine the whole chest i.e. interiorly, laterally and posteriorly.



#### **CAUSES:**

#### **DECREASED:**

- Pneumothorax
- Pleural effusion
- COPD

#### **INCREASED**:

- Pneumonia
- Pulmonary fibrosis
- Tumor


#### 4. APEX BEAT:

Localize the apex beat and note any shift i.e. towards right or left or in normal place.

#### **PERCUSSION**

Percuss all the lobes of lung and compare both sides. Follow the pattern as shown in the fig below.

Right lung has three lobes i.e. upper middle and lower lobe and left lung has two lobes upper and lower. Left middle lobe is called lingula.



<u>Note:</u> while percussing it is ideal to percuss and auscultate in the intercostal spaces in lean person and in any region in obese patient.

Start from the apex i.e. percuss the apices of the lung in supraclavicular fossa and then from 2<sup>nd</sup> to 6<sup>th</sup> intercostal spaces in front, and from 4<sup>th</sup> to 7<sup>th</sup> on lateral sides. Resonant sounds= normal (loud and low pitched) **Hyper resonant** =pneumothorax, asthma or emphysema (very loud, low pitch and longer in duration) Tympanic= empty stomach (over gas filled spaces. loud and high pitch)

Impaired note= fibrosis

Dull note= collapse + consolidation + Obesity (pneumonia, hemothorax)


**Stony dull**= pleural effusion and empyema

#### **IMPORTANT:**

Always use the one or two fingers and movement should entirely at wrist and not at the elbow. And always compare both sides at every step.

#### **AUSCULTATION**

Steps are the same as percussion. Auscultate the apices of the lung and then from  $2^{nd}$  to  $6^{th}$  intercostal space anteriorly. From axilla to  $8^{th}$  rib laterally and ask to breath.



Эдее-33

Asses the type of breathing.

**Normal**= vesicular breathing **Harsh** = bronchial breathing

Added sounds (wheezes, crackles etc.)

➤ **VOCAL RESONANCE:** similar to vocal fremitus but is heard with stethoscope and ask the patient to say 3, 3, 3(teen teen teen) and note resonance.

Causes: same as vocal fremitus

Vesicular Breathing	Bronchial Breathing (tracheal)
Inspiration > expiration	Expiration > inspiration
No pause b/w Inspiration expiration	& Pause in between
Rustling sound (soft light sound)	Blowing character/ high pitched
Normal sound	Seen in pneumonia, heart failure, COPD
<b>CRACKLES</b> (Clicking, rattl like the take tak of clock)	ing or cracking soundi.e.
FINE CRACKLES	COARSE CRACKLES
Inspiration > expiration	Expiration > inspiration

No pause b/w inspiration and expiration	Pause in between
Rustling sound (soft light sound)	Blowing character (sound)
Low pitched	High pitched
e.g. pulmonary fibrosis, CCF	e.g. Bronchiectasis

**Note:** Crackles that do not clear after cough usually indicate pulmonary edema or fluid in alveoli due to ARDS or heart failure.

Crackles that partially clear or change after cough may show bronchiectasis ....

**WHEEZES:** "High pitched, coarse whistling sound heard during breathing".

- Wheezes during expiratory phase shows bronchiolar disease.
- ii. Wheezes during **inspiratory phase** shows stenosis caused by tumor, foreign bodies or scarring. Also occurs in hypersensitivity pneumonitis.

#### **STRIDOR: (MUSICAL BREATHING)**

Special type of wheeze, harsh, high pitched and vibrating sound

• **Inspiratory**, which occurs when breathing in, and indicates a blockage above the vocal cords. E.g.

- croup, epiglottitis, laryngomalacia, vocal cord paralysis,
- **Expiratory**, which occurs when breathing out, and indicates a blockage in the windpipe. E.g. tracheal stenosis, anaphylaxis, bacterial tracheitis etc
- Biphasic, which occurs when a person breathes in and out, and indicates narrow cartilage directly below the vocal cords. e.g. Foreign body, hemangioma, mass or tumor.

Pleural friction rub	Crackles
ls biphasic	Either inspiratory or expiratory or both
Usually localized to a small area	Widespread
No change after coughing	May clear after coughing
Often palpable	Usually nonpalpable
Pressure of stethoscope intensify the sounds	No effect
Associated with pleuritic chest pain or local tenderness	No pain or tenderness

- Properly cover the patient
- Thank the patient

#### **IF THE COMMAND IS:**

## **EXAMINE THE PATIENT FROM THE BACK FOR CHEST EXAMINATION**

Same steps i.e. RIPE

<u>Position:</u> while examining back of the patient take off the shirt and make the position which is most suitable to you. Ask the **patient to sit with arms crossed anteriorly.** 

#### **INSPECTION:**

Same as that from front, also look for kyphosis, scoliosis or Kyphoscoliosis etc.

#### **PALPATION:**

Always ask for tenderness and pain.

While palpating apices of the lungs place the four fingers from behind in the supraclavicular fossa and thumb stretched behind the neck.

Rest of the palpation is same as that of palpation from front but no apex beat.

#### **PERCUSSION & AUSCULTATION:**

Same as from the front side.

- Cover AND thank the patient.
- NOTE: IF THE COMMAND IS: EXAMINE THE PATIENT FOR THE CHEST EXAMINATION THEN PERFORM ALL THE STEPS i.e. BOTH FROM THE FRONT AND BACK ALONG WITH RELEVANT

# **SECTION 2: FLUID ANALYSIS**

- 1. CEREBROSPINAL FLUID
- 2. ASCITIC FLUID
- 3. PLEURITIC FLUID
- 4. URINALYSIS

# **CEREBROSPINAL FLUID (CSF) ANALYSIS:**

CSF is a clear, colorless body fluid found in brain and spinal cord produced by specialized ependymal cells in choroid plexus of the ventricles of the brain.

CSF analysis is a group of tests that evaluate substances in CSF in order to diagnose conditions affecting the brain and spinal cord (CNS).

## TOTAL VOLUME = 100-150 ml DAILY SECRETION = 400-450 ml

Diseases are widely divided into four categories;

- Infectious disease: meningitis, encephalitis
- **Bleeding** with in the skull or brain
- Autoimmune disorders; GBS, MS, sarcoidosis
  - Tumors of CNS

#### PHYSICAL PROPERTIES OF CSF:

**AMOUNT:** Normally suitable amount for CSF analysis = **2-5ml** 

#### PRESSURE:

Measurement of pressure during collection i.e. during starting and closing the collection.

**Normal pressure** = 50-250 mm of water

Increased = tumors, infection, hydrocephalous or bleeding (SAH)

**Decreased** = dehydration, shock or leakage of CSF (sinus fracture)

**COLOUR:** Normally CSF is clear and colorless.

Color	Cloudy	Clear	Straw/cloudy /yellow	Blood stained(red)	Clear
Presentin	Bacterial meningitis	Viral meningitis	Tuberculosis meningitis	Subarachnoid hemorrhage or trauma during procedure.	Multiple sclerosis

#### **CHEMICAL TESTS:**

#### **CSF GLUCOSE:**

- Normally =  $2/3^{rd}$  of blood glucose
- **Decreased** (bacteria usually eat sugar so glucose decreases in) = bacterial meningitis (even up to 0 in severe cases) and somewhat in tuberculous meningitis below 40mg/dl
  - **Normal** = SAH , viral, MS

#### **CSF PROTEIN:**

- Normal = < 0.45g/L (20-45mg/dl) adults
- In newborn = 120-160 mg/dl are normal
  - Increased = increase in all conditions except in viral and MS which may be normal/increased
- CSF LACTIC ACID :

Increased = bacterial and fungal meningitis Normal in viral meningitis

• CSF LDH:

Normal In = viral meningitis Increased in = bacterial meningitis, leukemia, stroke

#### **MICROSCOPIC TESTS:**

- RBCs: normally not present or 0-2 are normal
- If present more than 2 then it shows SAH or trauma during procedure. Also increase in herpes virus due to vascular involvement
- **WBCs**: **normal** = 0-5 cells with 60 to 70 % lymphocytes (adults)

**Neonates** = up to 30 cells are normal **Increased** in infections etc.

- Bacterial meningitis = from 100-60000+ with neutrophils predominant
- **Viral meningitis** = 10-100 with lymphocytes predominant
- Tuberculous meningitis = 10-500 first neutrophils predominate then lymphocytes
- Fungal meningitis = monocytes predominate

#### **TESTS FOR INFECTIONS:**

- CSF Gram Staining and C/S
- PCR For Virus Detection

## **SUMMARY**

			SI			
CONDITIONS	NORMAL	BACTERIAL MENINGITIS	TUBERCULOUS MENINGITIS	VIRAL MENINGITIS	SAH	FUNGAL
Pressure	50-250 mmH20	Normal /increase d	Normal/ increased	normal	Normal	Increased
Colour	Clear	Cloudy(depends on microorganism numbers)	Straw/yellow	clear	Reddish pink	1
Glucose mg/dl	2/3 <sup>rd</sup> of Glucose	Decreased even up to zero	Decreased up to 35mg/dl	normal	Normal	Dec
Proteins mg/dl	20-45mg/dl in adults in newborn 120-160 mg/dl	100-500	100-3000	20-100	20-100	High

Page-41

Races	0-2 cells	:	:		Increased	
Leukocytes	0-5=adults Up to 30 =neonates	100-1 lac mostly neutrophils	10-500 mostly lymphocytes	10-150 mostly lymphocytes		Lymphocytes predominate
ГОН	-	Increased	Increased	normal		
ACTIC ACID	:	Increased	Increased	normal		Increased

# **SECTION 3**

- ABGs
- DRUGS
- POISONS

# **Arterial Blood Gases (ABGs):**

#### **INDICATIONS:**

- Respiratory failure
- Cardiac failure
- Renal failure
- Liver failure
- Hyperglycemic state associated with DM
- Sepsis
- Burns
- Poison/toxins
- Ventilated patient

#### **CONTRAINDICATIONS:**

#### **ABSOLUTE:**

- Abnormal Allen test (no collateral blood flow to the hand)
- Local infection
- Distorted anatomy at puncture site
- AV fistulas
- Severe peripheral vascular disease of limb involved

#### RELATIVE:

- Severe coagulopathy
- Anticoagulant therapy i.e. warfarin
  - Use of thrombolytic agents i.e. streptokinase etc.

#### In ABGs we see the

- PH
- PaCO<sub>2</sub>
- Bicarbonate
- Anion gap

## • Alveolar-arterial oxygen gradient (A-a)

• PaO<sub>2</sub>

			1
	Normal	Acidotic	Alkalotic
PH	7.35-7.45	< 7.35	>7.45
Paco2	35-45 mmHg	>45	<35
	(4.7-6.0 kPa)		
Нсо3	22-26mEq/L	<22	>26
Pa02	80-100 mmHg	-	-
Base excess (surrogate marker of metabolic acidosis or alkalosis)	-2 to +2 mmol/L	-2mmol/L shows that there is lower than normal amount of HcO3 in blood due to primary metabolic acidosis or compensated respiratory alkalosis	+2mmol/L (either due to primary metabolic alkalosis or compensated respiratory acidosis)
435	3211503		


Page-45


## **SECTION 4**

- **INSTRUMENTS**
- Hasairs oscii, Qidiridii on X-RAYS

## **INSTRUMENTS**

- NG TUBE
- FOLEY'S CATHETER
- IV CANNULA
- INSULIN SYRINGE
- LP NEEDLE
- ETT
- PEAK FLOW METER
- INHALERS (MDI + DISKUS)
- SPACER
- VENTURI MASK
- OROPHYRANGEAL AIRWAY
- NASOPHARYNGEALAIRWAY
- INTEROSSEUS CANNULA
- TRUCUT BIOPSY NEEDLE
- BLOOD SAMPLETUBES
- BONEMARROWBIOPSYNEEDLE (jamshidi needle)
- PLEURAL BIOPSY NEEDLE (Abram's needle)
- CENTRAL VENOUS LINE
- MONOPTY GUN
- AMBO BAG
- NEONATAL BP CUFF





## NG TUBE (NASO-GASTRIC TUBE)

**TYPES:** 1.Ryle's tube

2. Levine's tube

## **INDICATIONS:**

- Feeding
- Gastric lavage as in poisoning etc.
- Administration of medications
- Aspiration of gastric content for diagnostic purpose

- Stomach decompression as in obstruction etc.
- Enteral nutrition
- Prolonged ileus
- UGI bleeding confirmation

#### **CONTRAINDICATIONS:**

- Midface injury and basilar skullfracture (b) Hx of gastric bypass
- Hxof acid/alkali ingestion (d) Esophageal strictures
- Coagulopathy


#### **COMPLICATIONS:**

- a) Bleeding (b) Perforation (c) Wrong placement
- d) Pneumothorax (e) aspiration
- (f) vomiting/retching

SIZES: 12-18 French (fr) in adults, In neonates

8-10fr,in infants and children less than 5 years =

8-12 fr





## **URINARY CATHETHER (FOLEY'S catheter)**

i. TYPES: Materials: silicon and rubber

Procedure: double lumen and triple

and others shown in fig.

#### **INDICATIONS:** (

THERAPEUTIC	DIAGNOSTIC
1. Comatose patient	Renal function tests
2. Urinary incontinence	Residual volume
3. Urinary retention	Radiographic

## contrast studies

- 4. Intravesical chemotherapy
- 5. In all perineal operations
- 6. In Cerebrovascular accident

## **CONTRAINDICATIONS:**

1.	Acute	2.	Blood at
	urethritis		meatus
3.	Urethral	4.	Scrotal
	trauma		hematoma
5.	High		
	riding		
	prostrate		

## **COMPLICATIONS:**

Trauma & hemorrhage	UTI
False passage , tube blockage	Fistula formation

# Page-52

# **CHEST X-RAYS (CXR)**

IDENTIFICATION: First of all look for

1) Name (2) Date

Type of film:

Position: PA view, AP view, lateral view,

 lateral decubitus (for sub pleural effusion and when pleural effusion is less or doubtful)

• **Direction**: upright etc

PA VI	EW	AP VIEW
•	Apex of lung above the clavicle due to abduction of arm	Apex of the lung below the clavicle
•	Body of scapula visible (scapula move out)	Not visible
•	Heart size is normal	Comparatively appears bigger
•	Chin not visible	Chin visible
•	Posterior ribs appears more superficial as compared to anterior ribs	Not so
•	Not marked as PA	Usually marked as AP

### Technical quality: "RIP"

**R= Rotation** (the clavicular ends must be at equal distance from the spinous process (vertebral spine shadow) (tear drop)

- **I= Inspiration** (the CXR must be taken in full inspiration and 6<sup>th</sup> anterior rib must pass through the mid of Right .hemi diaphragm)
- P= Penetration/exposure (draw a horizontal line through the most concave part of cardiac shadow.

### **NORMAL EXPOSURE:**

Vertebrae should be visible above and disappear as we move down the film i.e. not all vertebrae are visible means Visible in cervical region only, slightly visible in chest region and disappears below the diaphragm.

#### OVER-EXPOSURE:

If whole of vertebral column is visible.

# Look at the big picture

### **MEDIASTINUM:**

## **Divisions:**

Superior and inferior (divided by horizontal line passing through the angle of Louis)

Anterior, middle, posterior

## **SUPERIOR MEDIASTINUM:**

Trachea, Esophagus, Thymus (up to 8 to 18 months of age)

#### INFERIOR MEDIASTINUM:

Heart, Lungs, Wide mediastinal shadow = **HOTS** 

H= Histiocytosis

O= Onco i.e. carcinoma lymphoma etc.

T = TB, thymus

S= Silicosis

#### **HEART:**

#### LOCATION:

- 1/3<sup>rd</sup> to the right
- 2/3<sup>rd</sup> to the left (normal)

#### **BORDERS:**

- **Right border**: by right atrium and IVC
- Left border: left ventricle and some of left atrium
- Inferior border: left and right ventricle
- **Superior border:** right and left atrium and the great vessels.
- SIZE: Normal cardiothoracic ratio (CTR) = < 55%</li>

Cardiomegaly = >55% of CTR


### **Transthoracic diameter (TTD) or CTR:**

- Draw a horizontal line passing just above the dome of the right diaphragm. (label it as A)
- Draw a perpendicular line in the midline

- Draw a horizontal line from the most concave part of the heart on the right side toward the midline. (label it as B)
- Draw the same line but from the left side of the heart. (label it as C)

#### **FORMULA**

 $TTD = (B+C/A) \times 100$ 



# SECTION 5 PROCEDURES

- Nebulization
- Neonatal resuscitation
- IV Cannula
- NG tube insertion
- Female Catheterization
- Male Catheterization
- CPR
- > COUNSELLING
- **EPI SCHEDULE**
- > ADMISSION CRITERIA FOR NURSERY
- GYNAE & SURGERY INSTRUMENTS

# **PROCEDURES**

## **NEBULIZATION:**

## **Apparatus:**

- 1) Mask
- 2) Chamber
- 3) Tubing
- 4) Nebulization machine



#### Dose:

- 0.5 ml Ventolin + 0.5 ml normal saline (If child is < 1 yr)</li>
- 1 ml Ventolin + 1 ml normal saline (If child is > 1 yr)
- Standard dose: 1:1 but, in clinical practice 2.5ml normal saline is used for 0.5 ml Ventolin (0.5ml Ventolin= 16 drops)
- Normal saline is hypertonic for airway=> oozes out fluids
- In bronchiolitis=> only normal saline is used for nebulization
- Ventolin nebulization include 3 to 4 nebulization's at one time with a break of 10-15 minutes
- Anticholinergics are used in a child with congenital heart disease in place of Ventolin.
- Betamethasone is used for nebulization when there is severe inflammation

## **COUNSELLING**

## **MEDICAL COUNSELLING STEPS**

- Introduce yourself and confirm the disease and patient (ask his name)
- Take permission for counselling (in front of the attendant or in separate room)
- Ask the pt. about his/her education, and knowledge about his/her disease?
- Ask the pt. whether he/she is counselled before or not?
- Ask the pt. complaints very briefly?
- Describe the disease.
- Pathology -> signs and symptoms
- Investigations
- Diagnosis
- Treatment modalities available
- Describe/explain drugs, duration, adverse effects.
- Compliance to drugs and follow up.
- You can ask any question if you have
- Say thanks to patient.

- **♦** السلام وعليكم!!!
- 1. میرانام ڈاکٹر XYZ ہے۔اور میں ڈاکٹروں کے اس ٹیم کا حصہ ہوں جو آپنے نیچے کاعلاج کروار ہی ہے۔
  - 2. آپ اس مریض کے والد صاحب ہے؟ جس کوید بیاری ہے۔
- 3. میں آپ کے بیٹا/بیٹی کے بیاری کے بارے میں کچھ باتیں آپ سے کرنا چاہتا ہوں توسب کے سامنے آپ سننا پیند کریں گے یا کیلے میں۔
- 4. آپ کی تعلیم کتنی ہے۔ اور آپ اپنے مریض کے بیاری کے بارے میں کیا جانتے ہیں۔
- 5. اب میں آپ کو آپ کے مریض کے بارے میں بناؤں گا۔ آپ کے مریض کو جو بیاری ہے اسکا مطلب ہے۔ " بخار کے ساتھ جھٹکے لگنا بغیر دماغی انفیکشن کے "انگلش میں اسے Febrile fits کہتے ہیں۔
- ✓ اس میں بچوں کو بخار کے ساتھ جھنگے کا خطرہ ہوتاہے۔ ہر 100 میں تقریباً 5-2 بچوں میں سے ہوتا ہے۔ دماغی طور پر عموماً بالکل ٹھیک ہوتے ہیں۔
- ✓ عموماً بخار کے ساتھ جھٹکے 18 سے 24 ماہ کی عمر میں پڑ سکتے ہیں۔ لیکن ضروری نہیں کہ یہ 24 کے در میان میں ہوبلکہ 6 ماہ سے 5 سال کے در میان آسکتے ہیں۔

- 2.Complex 1.Simple بخار میں جھٹے دو طرح کے ہوتے ہیں۔
- 1. یہ جو Simple ہوتے ہیں سب سے زیادہ پائی جانے والی قتم ہے۔ اور صرف ایک 1-15 منٹ کا جھٹکا ہوتا ہے۔ اور پورے جسم پر ہوتا ہے۔ (Generalized)
- 24. Complex جوہ جسم کے ایک خاص جھے (Focal) میں مسلسل 24 گفتے ہوتا ہے۔ یا بار بار آتی ہے۔ بچہ کو عارضی طور پر بازو یا legl میں جھکے کے بعد کمزوری محسوس ہوتے ہیں۔
- Febrile Fits بین ـ 1 1 1 2 مختلف اسباب (Causes) بین ـ 1 4 Family Hx 3 (MMR )Immunization 2 4 Family Hx 3 (MMR ) المسيث کی خرابی 5 Malaria
  - ♣ Febrile Fits ڪعلامات پچھ ايسے ہوتے ہیں۔ بخار کے پہلے دن ہی جھڻکا آتا ہے جو اس بات کو ظاہر کرتا ہے کہ بچہ بیار ہے۔ زیادہ تر جھٹکا اس وقت آتا ہے ج5 °C) 102.2 °F سے زیادہ ہو جائے۔
    - 💠 مندرجه ذیل بچول میں بیہ بار بار ہوسکتا ہے۔
      - تقریباً 30-35 فیصد میں ایبا ہو سکتا ہے۔
        - 1. 15 ماہ سے کم عمر میں پہلا جھٹکاآیا ہو۔
          - 2. باربار بخار ہونا۔

- Family Hx .3
- 4. جھٹکے سے پہلے Low grade fever ہونا۔
  - 5. اگر بخار شروع ہوتے ہی جھٹکے لگے ہو۔
- ب اگر کسی بچے میں مندرجہ ذیل وجوہات میں سے دویادوسے زیادہ موجود ہوتو مرگی (Epilepsy) کے 10 فیصد حیانس آنے والے میں ہوسکتے ہیں۔
  - 1. اگر جھٹکوں کا دورانیہ 15 منٹ یااس سے زیادہ ہو (Complex)
    - 2. اگر جھنگے ایک سے زیادہ مرتبہ لگے ہو۔
    - 3. اگر جھکے جسم کے صرف ایک تھے میں ہوئے ہو۔
      - 4. اگر بچه ذہنی طور پر ٹھیک نہ ہو۔
  - پہلی مرتبہ جھٹکے لگنے کی صورت ڈاکٹر مریض کے پچھ ٹسٹ کریں گے جس
     سے مرض تشخیص میں مدو ملے گی۔
    - ♦ علاج:-
    - ❖ علان: -﴿ گھر پر علاج: - 1 - تیز بخار کی صورت میں بیچ کے کیڑے کم کر دے۔
      - 2۔ گرمیوں کے موسم میں اسے ٹھنڈی جگہ پر رکھیں۔
        - 3\_ بخار والاشربت م 6-4 گھنٹے بعد بلائیں۔


4- اگر کا جھٹکے کا دورانیہ 5 منٹ سے زیادہ ہو۔ تو پاخانے کے راستے سے
(Suppository) Valium) کی بتی اسے دیں یا فورا ہمپتال لے جائیں۔
5- اگر جھٹکوں کا دورانیہ 10 منٹ سے زیادہ ہو تو فورا ہمپتال لے جائیں۔
6- عموماً کمبے عرصے کے لئے دوائی ایسے جھٹکوں میں زیادہ دیر تک فائدہ نہیں
دیتی۔ تاہم بخار شروع ہوتے ہی Valium کی بتی پاخانے کے راستے صبح وشام
8- عموماً کم بخار شروع ہوتے ہی کا کا کہ بتی باخانے کے راستے صبح وشام

■ مزید کچھ معلومات در کار ہو تو مجھ سے پوچھ سکتے ہیں۔

# **INSTRUMENTS GYNAE AND SURGERY**









Hassains

## **BOOKS AVAILABLE AT:**

- → Swat Book House and Publications
  Opposite Saidu Medical College, Saidu Sharif Swat, KPK
  Cell# 0346-9049087, Ph# 0946-726817
- → London Book Bank, Near Ayub Medical College Abbottabad, KPK Ph# 0992-381157, Cell# 0321-9802166
- **→** Prince Bookshop

KTH Underpass, Phase 3 Chowk Hayatabad and Sadar Peshawar, KPK Contact# 0333-5556447, 091-5701086, 0333-9111926

- → **Khyber Bookshop,** Khyber Medical College Peshawar Safeer Cell# 0315-9739220, 0344-6403138
- → Noble Book Bank, Near STH Swat Cell# 0300-5746553
- → Noble Book Bank, Near Ayub Medical College Abbottabad, KPK Cell# 0300-9110858, 0992-383116
- → New Prince Bookshop, Dera Ismail Khan, KPK Cell# 0345-9889346
- → Prince Book Depot, Near Rawalpindi Medical College, Rawalpindi, Punjab Phone# 051-5502300, Cell# 0300-5121041
- → Global Book Bank, KDA Kohat, KPK Cell# 0333-9498473

- → Student Book Agency, Coffee Shop Peshawar, KPK Cell# 0312-9191998, 0334-9008093
- → Marwat Book Bank, Bannu, KPK Rooh-ul-Amin Cell# 0333-9847238
- → Rasheed Medical Books, Malakand Chowk, New Adda Mardan, KPK
  Cell# 0313-7810310, 0333-9890490